

Quadratics - 2
- Answers

Exercise A

1. Discriminant = $b^2 - 4ac$

$$= 5^2 - 4(1)(2)$$

$$= 25 - 8$$

$$= \underline{17}$$

2. Discriminant = $b^2 - 4ac$

$$= 4^2 - 4(2)(1)$$

$$= 16 - 8$$

$$= \underline{8}$$

3. Discriminant = $b^2 - 4ac$

$$= (-3)^2 - 4(4)(5)$$

$$= 9 - 80$$

$$= -71$$

4. Discriminant = 4

5. Discriminant = 56

6. Discriminant = 49

Exercise B

1.
$$b^2 - 4ac = 4^2 - 4(1)(3)$$

$$= 16 - 12$$

$$= 4 > 0$$

∴ Two distinct real roots.

2.
$$b^2 - 4ac = (-6)^2 - 4(2)(20)$$

$$= 36 - 160$$

$$= -124 < 0$$

∴ No real roots.

3.
$$x^2 + 6x + 9 = 0$$

$$b^2 - 4ac = 6^2 - 4(1)(9)$$

$$= 0$$

∴ One repeated real root.

4.
$$b^2 - 4ac = (-1)^2 - 4(3)(-13)$$

$$= 157 > 0$$

∴ Two distinct real roots.

5.
$$b^2 - 4ac = (-4)^2 - 4(1)(16)$$

$$= 16 - 64$$

$$= -48 < 0$$

∴ No real roots.

6.
$$b^2 - 4ac = (-2)^2 - 4(-1)(-10) = -36 < 0$$

$$∴ \text{No real roots.}$$

Exercise C

1. Since the equation has repeated real roots,

$$b^2 - 4ac = 0$$

$$(-6)^2 - 4(2)(k) = 0$$

$$36 = 8k$$

$$k = \underline{\underline{4.5}}$$

2.

(a) Discriminant = $b^2 - 4ac$
 $= (-4)^2 - 4(k)(k)$
 $= \underline{\underline{16 - 4k^2}}$

(b) Since there are equal roots, (repeated real roots),

$$b^2 - 4ac = 0$$

$$\therefore 16 - 4k^2 = 0$$

$$16 = 4k^2$$

$$k^2 = 4$$

$$k = \underline{\underline{\pm 2}}$$

3.

(a) Discriminant = $b^2 - 4ac$
 $= 7^2 - 4(-2)(3)$
 $= 49 + 24$
 $= 73$

Since the discriminant is greater than 0, there are two distinct real roots.

4.(a) $a=k, b=4, c=5-k$

Since there are two distinct real roots,

$$b^2 - 4ac > 0$$

$$4^2 - 4k(5-k) > 0$$

$$16 - 20k + 4k^2 > 0$$

($\div 4$)

$$4 - 5k + k^2 > 0$$

$$\therefore k^2 - 5k + 4 > 0$$




(b) $k^2 - 5k + 4 > 0$

To solve, sketch the graph of $y = k^2 - 5k + 4$.

Critical values: $k^2 - 5k + 4 = 0$

$$(k-4)(k-1) = 0$$

$$k = 4, 1$$



$$\therefore k < 1 \text{ or } k > 4$$



(You do not have to find the turning points unless you are asked to find...)

Exercise D


1. x -intercepts:

$$x^2 - 8x + 12 = 0$$

$$(x-6)(x-2) = 0$$


$$x = 6, 2$$

y -intercept = 12




2. $(x+5)(x+1) = 0$

$$x = -5, -1$$



3. $(x-7)(x+2) = 0$

$$x = 7, -2$$




4. $x^2 - 4x + 1 = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{4 \pm \sqrt{16 - 4(1)(1)}}{2(1)}$$


$$x = 2 + \sqrt{3}, 2 - \sqrt{3}$$



5. $x^2 + 10x - 3 = 0$

$$x = \frac{-10 \pm \sqrt{100 - 4(1)(-3)}}{2(1)}$$


$$x = -5 + 2\sqrt{7}, -5 - 2\sqrt{7}$$



6. $x^2 - 5x = 0$

$x(x - 5) = 0$


$x = 0, 5$



9. $2x^2 - 12x = 0$

$2x(x - 6) = 0$


$x = 0, 6$



7. $2x^2 - x - 3 = 0$

$(2x - 3)(x + 1) = 0$


$x = \frac{3}{2}, -1$



10. $x^2 - 9 = 0$

$x^2 = 9$


$x = \pm 3$



8. $3x^2 - 4x - 4 = 0$


$(x - 2)(3x + 2) = 0$

$x = 2, -2/3$



11. $x^2 - 25 = 0$


$x = \pm 5$



12. $2x^2 - 15 = 0$

$$2x^2 = 15$$

$$x = \pm \sqrt{\frac{15}{2}}$$



Turning point:

$$y = x^2 + 2x + 10$$


$$= (x+1)^2 - 1 + 10$$

$$= (x+1)^2 + 9$$

$$x+1=0 \quad y=+9$$

$$x = -1$$

$$(-1, 9)$$



13. x -intercepts:

$$x^2 + 2x + 10 = 0$$

$$x = \frac{-2 \pm \sqrt{4 - 4(1)(10)}}{2(1)}$$

$$x = \frac{-2 \pm \sqrt{-36}}{2}$$

No real roots.

\therefore No x -intercepts.

y -intercept = 10.

If there are no x -intercepts, you should find out the turning points to correctly sketch the graph, because it can tell you whether the turning point is to the left or right of the y -axis.

14.

$$b^2 - 4ac = (-1)^2 - 4(1)(5)$$

$$= -19 < 0$$

$\therefore x^2 - x + 5 = 0$ has no real roots. Hence the graph has no x -intercepts

y -intercept = 5


Turning point:


$$y = x^2 - x + 5$$

$$= \left(x - \frac{1}{2}\right)^2 - \frac{1}{4} + 5$$

$$= \left(x - \frac{1}{2}\right)^2 + \frac{19}{4}$$

$$\left(\frac{1}{2}, +\frac{19}{4}\right)$$



15. Shape: 

$$b^2 - 4ac = 4^2 - 4(-1)(-20)$$

$$= -64 < 0$$

∴ No x-intercepts.

y-intercept = -20

Turning point:

$$y = -x^2 + 4x - 20$$

$$= -1[x^2 - 4x] - 20$$

$$= -1[(x-2)^2 - 4] - 20$$

$$= -(x-2)^2 + 4 - 20$$


$$= -(x-2)^2 - 16$$

$$x-2=0$$

$$x=2$$

$$y = -16$$

$$(2, -16)$$



16. $b^2 - 4ac = 3^2 - 4(-2)(-15)$

$$= -111 < 0$$

∴ No x intercepts.

y-intercept = -15

Shape: 

Turning point:

$$y = -2x^2 + 3x - 15$$


$$= -2\left[x^2 - \frac{3}{2}x\right] - 15$$

$$= -2\left[\left(x - \frac{3}{4}\right)^2 - \frac{9}{16}\right] - 15$$

$$= -2\left(x - \frac{3}{4}\right)^2 + \frac{9}{8} - 15$$

$$= -2\left(x - \frac{3}{4}\right)^2 - \frac{111}{8}$$

$$\left(\frac{3}{4}, -\frac{111}{8}\right)$$



Exercise E

1. Let $y = x^2$
 Then, $y^2 = x^2 \times x^2 = x^4$
 $\therefore y^2 - 5y + 4 = 0$
 $(y-1)(y-4) = 0$
 $y = 1$ or $y = 4$
 $x^2 = 1$ or $x^2 = 4$
 $x = \pm 1$ or $x = \pm 2$

2. Let $y = x^2$
 $y^2 - 3y - 10 = 0$
 $(y-5)(y+2) = 0$
 $y = 5$ or $y = -2$
 $x^2 = 5$ or $x^2 = -2$
 $x = \pm 5$ No \downarrow real roots.

3. Let $y = x^3$
 Then, $y^2 = x^3 \times x^3 = x^6$
 $\therefore 2y^2 + 9y + 4 = 0$
 $2y^2 + 8y + y + 4 = 0$
 $2y(y+4) + 1(y+4) = 0$
 $(y+4)(2y+1) = 0$
 $y = -4$ or $y = -\frac{1}{2}$
 $x^3 = -4$ or $x^3 = -\frac{1}{2}$

$x = \sqrt[3]{-4}$ or $x = \sqrt[3]{-\frac{1}{2}}$

4. Let $y = x^{\frac{1}{4}}$
 Then $y^2 = x^{\frac{1}{4}} \times x^{\frac{1}{4}} = x^{\frac{1}{2}}$
 $\therefore 2y^2 - y - 3 = 0$
 $2y^2 - 3y + 2y - 3 = 0$
 $y(2y-3) + 1(2y-3) = 0$
 $(2y-3)(y+1) = 0$
 $y = \frac{3}{2}$ or $y = -1$

$x^{\frac{1}{4}} = \frac{3}{2}$ or $x^{\frac{1}{4}} = -1$
 $x = \left(\frac{3}{2}\right)^4$ or $x = (-1)^4$
 $x = \frac{81}{16}$ or $x = 1$

5. Let $y = x^{\frac{1}{6}}$
 Then $y^2 = x^{\frac{1}{6}} \times x^{\frac{1}{6}} = x^{\frac{1}{3}}$
 $\therefore 3y^2 - 9y = 0$
 $3y(y-3) = 0$
 $y = 0$ or $y = 3$
 $x^{\frac{1}{3}} = 0$ or $x^{\frac{1}{3}} = 3$
 $x = 0^3$ or $x = 3^3$
 $x = 0$ or $x = 27$