1.

Magnetic Fields 3

a)	State Faraday's law of electromagnetic induction.
	[1]

(b) Fig. 5.1 shows a magnet being moved towards the centre of a flat coil.





Fig. 5.1

A current is induced in the coil. Use ideas about energy conservation to state and explain the polarity of the face of the coil nearer the magnet.

[1]

(c) Fig. 5.2 shows the magnetic field from the north pole of a vertically held bar magnet.



(i) A small flat coil is placed at A. The coil is moved downwards from position A to position B. The plane of the coil remains horizontal between these two positions. Explain why there is no induced e.m.f. across the ends of the coil.

[1]

(ii) Fig. 5.3 is a graph showing how the magnetic flux density *B* varies along the horizontal line **XY** in Fig. 5.2.




Fig. 5.3




Fig. 5.4

The same small flat coil from (i) is moved at a constant speed from **X** to **Y**. The plane of the coil remains horizontal between **X** and **Y**.


On the axis provided in Fig. 5.4, sketch a graph to show the variation of the induced e.m.f. *E* across the ends of the coil with distance from **X**. [3]

2.	
(a)	State Lenz's law

	••••
(2 mar)	


(b) Figure 4 shows two small, solid metal cylinders, P and Q.P is made from aluminium. Q is made from a steel alloy.

Figure 4



(b) (i)	The dimensions of P and Q are identical but Q has a greater mass than P . Explain what material property is responsible for this difference.
	(1 mark)
(b) (ii)	When $\bf P$ and $\bf Q$ are released from rest and allowed to fall freely through a vertical distance of 1.0 m, they each take 0.45 s to do so. Justify this time value and explain why the times are the same.
	(2 marks)

(c) The steel cylinder Q is a strong permanent magnet. P and Q are released separately from the top of a long, vertical copper tube so that they pass down the centre of the tube, as shown in **Figure 5**.




The time taken for Q to pass through the tube is much longer than that taken by P.

(c) (i)	Explain why you would expect an emf to be induced in the tube as ${f Q}$ passes through it.
	(2 marks)
(c) (ii)	State the consequences of this induced emf, and hence explain why ${\bf Q}$ takes longer than ${\bf P}$ to pass through the tube.
	(3 marks)
(d)	The copper tube is replaced by a tube of the same dimensions made from brass. The resistivity of brass is much greater than that of copper. Describe and explain how, if at all, the times taken by $\bf P$ and $\bf Q$ to pass through the tube would be affected.
	P :
	Q:
	(3 marks)

A uniform magnetic field is produced by mounting two flat magnets on a U-shaped iron frame, so that the north and south poles are facing, as shown in **Figure 3**. The flux density of the magnetic field is 45 mT and may be assumed to act only over the area of the pole faces, which measure 40 mm by 20 mm. This magnet arrangement rests on the pan of a top pan balance.

Figure 3



(a) A horizontal wire is placed in the centre of the magnetic field and aligned to make it perpendicular to the flux lines. When a current is passed through the wire, the balance reading increases by 1.4×10^{-3} kg.

Calculate the current in the wire.	
	(2 marks)

(b) The wire is disconnected from the current source and its ends are connected to a sensitive voltmeter. When the wire is moved rapidly, vertically upwards across the whole magnetic field, cutting all of the flux lines perpendicularly, the voltmeter gives a reading.

Calculate

(b)	(i)	the magnetic flux change experienced by the wire during its movement completely across the magnetic field,

taken for the wire to pass completely across the maded at constant speed, if the voltmeter reads 0.15 m	wire to pass completely across the magnetic field, assuming speed, if the voltmeter reads 0.15 mV.		
 	(5 marks)		
	(5 mar		

4.

A single-turn square coil of side $0.050\,\mathrm{m}$ is placed in a magnetic field of flux density B of magnitude $0.026\,\mathrm{T}$.

(a) The coil is placed in three different orientations to the field as shown in Fig. 6.1(a), (b) and (c).




Fig. 6.1

In Fig. 6.1(a), the plane of the coil is perpendicular to the field. In (b), it is at 45° to the field and in (c), it is parallel to the field. Calculate the value, giving a suitable unit, of the magnetic flux linking the coil for the position shown in

(i) Fig. 6.1(a)

magnetic flux =unit....... [3]

(ii) Fig. 6.1(b)

magnetic flux =unit....... [1]

(iii) Fig. 6.1(c).

magnetic flux =unit.......[1]

(b) The coil is rotated in the magnetic field to generate an e.m.f. across its ends. The graph of the variation of e.m.f. with time is shown in Fig. 6.2.




Fig. 6.2

(i)	On Fig. 6.2 mark, with an X, a point on the graph at a time when the flux linking	the)
	coil is a maximum.	[1]


(ii)	Give your reasoning for your choice of position X.		
	[2		

(iii) The rate of rotation of the coil is doubled. On Fig. 6.2 draw a graph showing at least two cycles of the e.m.f. now generated across the ends of the coil. [3]

5.

(a) Figure 5 shows two coils, P and Q, linked by an iron bar. Coil P is connected to a battery through a variable resistor and a switch S. Coil Q is connected to a centre-zero ammeter.


Figure 5



(a) (i)	Initially the variable resistor is set to its minimum resistance and ${\bf S}$ is open. Describe and explain what is observed on the ammeter when ${\bf S}$ is closed.	[3 marks]
(a) (ii)	With S still closed, the resistance of the variable resistor is suddenly increas Compare what is now observed on the ammeter with what was observed in Explain why this differs from what was observed in part (a)(i).	

(b) Figure 6 shows a 40-turn coil of cross-sectional area $3.6 \times 10^{-3} \, \mathrm{m}^2$ with its plane set at right angles to a uniform magnetic field of flux density $0.42 \, \mathrm{T}$.

Figure 6



(b) (i)	Calculate the magnitude of the magnetic flux linkage for the coil. State an appropriate unit for your answer.	[2 marks]
	flux linkage	unit
(b) (ii)	The coil is rotated through $90^{\rm o}$ in a time of $0.50~\rm s.$ Determine the mean emf in the coil.	[2 marks]
	mean emf	V