

Sequences 2 - Answers

1.

(a)	John; arithmetic series, $a = 60$, $d = 15$.		
	$60 + 75 + 90 = 225^*$ or $S_3 = \frac{3}{2}(120 + (3 - 1)(15)) = 225^*$	Finds and adds the first 3 terms or uses sum of 3 terms of an AP and obtains the printed answer, with no errors.	B1 *
	Beware: The 12 th term of the sequence is 225 also so look out for $60 + (12 - 1) \times 15 = 225$. This is B0.		
			[1]
(b)	$t_9 = 60 + (n - 1)15 = (£)180$	M1: Uses $60 + (n - 1)15$ with $n = 8$ or 9 A1: (£)180	M1 A1
	Listing: M1: Uses $a = 60$ and $d = 15$ to select the 8 th or 9 th term (allow arithmetic slips) A1: (£)180 (Special case (£)165 only scores M1A0)		
			[2]
(c)	$S_n = \frac{n}{2}(120 + (n - 1)(15))$ or $S_n = \frac{n}{2}(60 + 60 + (n - 1)(15))$	Uses correct formula for sum of n terms with $a = 60$ and $d = 15$ (must be a correct formula but ignore the value they use for n or could be in terms of n)	M1
	$S_n = \frac{12}{2}(120 + (12 - 1)(15))$	Correct numerical expression	A1
	$= (£)1710$	cao	A1
	Listing: M1: Uses $a = 60$ and $d = 15$ and finds the sum of at least 12 terms (allow arithmetic slips) A2: (£)1710		
			[3]
	(d)	$3375 = \frac{n}{2}(120 + (n - 1)(15))$	Uses correct formula for sum of n terms with $a = 60$, $d = 15$ and puts $= 3375$
	$6750 = 15n(8 + (n - 1)) \Rightarrow 15n^2 + 105n = 6750$	Correct three term quadratic. E.g. $6750 = 105n + 15n^2$, $3375 = \frac{15}{2}n^2 + \frac{105}{2}n$ This may be implied by equations such as $6750 = 15n(n + 7)$ or $3375 = \frac{15}{2}(n^2 + 7n)$	A1
	$n^2 + 7n = 25 \times 18^*$	Achieves the printed answer with no errors but must see the 450 or 450 in factorised form or e.g. 6750, 3375 in factorised form i.e. an intermediate step.	A1*
			[3]
(e)	$n = 18 \Rightarrow \text{Aged } 27$	M1: Attempts to solve the given quadratic or states $n = 18$ A1: Age = 27 or just 27	M1 A1
	Age = 27 only scores both marks (i.e. $n = 18$ need not be seen)		
	Note that (e) is not hence so allow valid attempts to solve the given equation for M1		
			[2]

2.

	(a) Use $n^{\text{th}} \text{ term} = a + (n-1)d$ with $d = 10$; $a = 150$ and $n = 8$, or $a = 160$ and $n = 7$, or $a = 170$ and $n = 6$: $= 150 + 7 \times 10$ or $160 + 6 \times 10$ or $170 + 5 \times 10$ $= 220^*$ (Or gives clear list – see note)	M1 A1*	(2)
Or	If answer 220 is assumed and $150 + (n-1)10 = 220$ or variation is solved for n = Then $n = 8$, so 2007 is the year (must conclude the year)	M1 A1*	(2)
	(b) Use $S_n = \frac{n}{2} \{2a + (n-1)d\}$ $= \frac{7}{2} (300 + 13 \times 10)$ $= 7 \times 430$ $= 3010$ Or $S_n = \frac{n}{2} \{a + l\}$ and $l = a + (n-1)d$ or $7(150 + 280)$	M1 A1 A1	(3)
	(c) Cost in year $n = 900 + (n-1) \times 20$ Sales in year $n = 150 + (n-1) \times 10$ Cost = 3 × Sales $\Rightarrow 900 + (n-1) \times 20 = 3 \times (150 + (n-1) \times 10)$ $900 - 20n + 20 = 450 + 30n - 30$ $500 = 50n$ $n = 10$ Year is 2009 As n is not defined they may work correctly from another base year to get the answer 2009 and their n may not equal 10. If doubtful – send to review.	M1 M1 M1 A1	(4)

3.

(a)	$600 = 200 + (N-1)20 \Rightarrow N = \dots$	Use of 600 with a correct formula in an attempt to find N . A correct formula could be implied by a correct answer.	M1
	$N = 21$	cso	A1
	Accept correct answer only.		
	$600 = 200 + 20N \Rightarrow N = 20$ is M0A0 (wrong formula) $\frac{600-200}{20} = 20 \therefore N = 21$ is M1A1 (correct formula implied)		
	Listing: All terms must be listed up to 600 and 21 correctly identified. A solution that scores 2 if fully correct and 0 otherwise.		
			(2)
(b)	Look for an AP first:		
	$S = \frac{21}{2} (2 \times 200 + 20 \times 20)$ or $\frac{21}{2} (200 + 600)$ or $S = \frac{20}{2} (2 \times 200 + 19 \times 20)$ or $\frac{20}{2} (200 + 580)$ (= 8400 or 7800)	M1: Use of correct sum formula with their integer $n = N$ or $N-1$ from part (a) where $3 < N < 52$ and $a = 200$ and $d = 20$. A1: Any correct un-simplified numerical expression with $n = 20$ or $n = 21$ (No follow through here)	M1A1

	Then for the constant terms:		
	$600 \times (52 - "N") (= 18600)$	M1: $600 \times k$ where k is an integer and $3 < k < 52$ A1: A correct un-simplified follow through expression with their k consistent with n so that $n + k = 52$	M1A1ft
	So total is 27000	Cao	A1
	Note that for the constant terms, they may correctly use an AP sum with $d = 0$.		
	There are no marks in (b) for just finding S_{52}		
			(5)
			[7]
	<p>If they obtain $N = 20$ in (a) (0/2) and then in (b) proceed with, $S = \frac{20}{2}(2 \times 200 + 19 \times 20) + 32 \times 600 = 7800 + 19\ 200 = 27\ 000$ allow them to 'recover' and score full marks in (b) Similarly If they obtain $N = 22$ in (a) (0/2) and then in (b) proceed with, $S = \frac{21}{2}(2 \times 200 + 20 \times 20) + 31 \times 600 = 8400 + 18\ 600 = 27\ 000$ allow them to 'recover' and score full marks in (b)</p>		

4.

(i).(a)	$U_3 = 4$	cao	B1
			(1)
(b)	$\sum_{n=1}^{n=20} U_n = 4 + 4 + 4 \dots + 4 \text{ or } 20 \times 4$	<p>For realising that all 20 terms are 4 and that the sum is required. Possible ways are $4+4+4 \dots +4$ or 20×4 or $\frac{1}{2} \times 20(2 \times 4 + 19 \times 0)$ or $\frac{1}{2} \times 20(4 + 4)$ (Use of a correct sum formula with $n = 20$, $a = 4$ and $d = 0$ or $n = 20$, $a = 4$ and $l = 4$)</p>	M1
	$= 80$	cao	A1
	Correct answer with no working scores M1A1		
			(2)
(ii)(a)	$V_3 = 3k, V_4 = 4k$	May score in (b) if clearly identified as V_3 and V_4	B1, B1
			(2)
(b)	$\sum_{n=1}^{n=5} V_n = k + 2k + 3k + 4k + 5k = 165$ <p>or</p> $\frac{1}{2} \times 5(2 \times k + 4 \times k) = 165$ <p>or</p> $\frac{1}{2} \times 5(k + 5k) = 165$	<p>Attempts V_5, adds their V_1, V_2, V_3, V_4, V_5 AND sets equal to 165 or Use of a correct sum formula with $a = k$, $d = k$ and $n = 5$ or $a = k$, $l = 5k$ and $n = 5$ AND sets equal to 165</p>	M1

$15k = 165 \Rightarrow k = ..$	Attempts to solve their linear equation in k having set the sum of their first 5 terms equal to 165. Solving $V_5 = 165$ scores no marks.	M1
$k = 11$	cao and cso	A1
		(3)

5.

	$a_1 = 4, a_{n+1} = 5 - ka_n, n \dots 1$	
(a)	$a_2 = 5 - ka_1 = 5 - 4k$ $a_3 = 5 - ka_2 = 5 - k(5 - 4k)$	<p>M1: Uses the recurrence relation correctly at least once. This may be implied by $a_2 = 5 - 4k$ or by the use of $a_3 = 5 - k$ (their a_2)</p> <p>A1: Two correct expressions – need not be simplified but must be seen in (a). Allow $a_2 = 5 - k4$ and $a_3 = 5 - 5k + k^2 4$ Isw if necessary for a_3.</p> <p>M1A1</p>
		[2]
(b)	$\sum_{r=1}^3 (1) = 1 + 1 + 1$	<p>Finds 1+1+1 or 3 somewhere in their solution (may be implied by e.g. $5 + 6 - 4k + 6 - 5k + 4k^2$). Note that $5 + 6 - 4k + 6 - 5k + 4k^2$ would score B1 and the M1 below.</p> <p>B1</p>
	$\sum_{r=1}^3 a_r = 4 + "5 - 4k" + "5 - 5k + 4k^2"$	<p>Adds 4 to their a_2 and their a_3 where a_2 and a_3 are functions of k. The statement as shown is sufficient.</p> <p>M1</p>
	$\sum_{r=1}^3 (1 + a_r) = 17 - 9k + 4k^2$	<p>Cao but condone ‘= 0’ after the expression</p> <p>A1</p>
	Allow full marks in (b) for correct answer only	
		[3]
(c)	500	cao
		B1
		[1]

6.

(a) $(S =) a + ar + \dots + ar^{n-1}$	“S=” not required.	Addition required.	B1
$(rS =) ar + ar^2 + \dots + ar^n$	“rS=” not required	(M: Multiply by r)	M1
$S(1 - r) = a(1 - r^n)$	$S = \frac{a(1 - r^n)}{1 - r}$	(M: Subtract and factorise) (*)	M1 A1 cso (4)
(b) $ar^{n-1} = 35000 \times 1.04^3 = 39400$	(M: Correct a and r , with $n = 3, 4$ or 5).		M1 A1 (2)
(c) $n = 20$	(Seen or implied)		B1

$S_{20} = \frac{35000(1-1.04^{20})}{(1-1.04)}$ <p>(M1: Needs <u>any</u> r value, $a = 35000$, $n = 19, 20$ or 21). (A1ft: ft from $n = 19$ or $n = 21$, but r must be 1.04). $= 1\,042\,000$</p>	M1 A1ft	
	A1	(4)
		10
<p>(a) B1: At least the 3 terms shown above, and no extra terms. A1: Requires a completely correct solution. <u>Alternative for the 2 M marks:</u> M1: Multiply numerator and denominator by $1 - r$. M1: Multiply out numerator convincingly, and factorise.</p> <p>(b) M1 can also be scored by a “year by year” method. <u>Answer only:</u> 39 400 scores full marks, 39 370 scores M1 A0.</p> <p>(c) M1 can also be scored by a “year by year” method, <u>with terms added</u>. In this case the B1 will be scored if the correct number of years is considered. <u>Answer only:</u> Special case: 1 042 000 scores 2 B marks, scored as 1, 0, 0, 1 (Other answers score no marks).</p> <p><u>Failure to round correctly in (b) and (c):</u> Penalise once only (first occurrence).</p>		

7.

(a)

	$\{S_n = \} a + ar + \dots + ar^{n-1}$ $\{rS_n = \} ar + ar^2 + \dots + ar^n$ $(1-r)S_n = a(1-r^n)$ $S_n = \frac{a(1-r^n)}{1-r} \quad (*)$	B1 M1 dM1 A1cso (4)
(b)	$a = 200, r = 2, n = 10, \quad S_{10} = \frac{200(1-2^{10})}{1-2}$ $= 204,600$	M1, A1 A1 (3)
(c)	$a = \frac{5}{6}, r = \frac{1}{3}$ $S_{\infty} = \frac{a}{1-r}, \quad S_{\infty} = \frac{\frac{5}{6}}{1-\frac{1}{3}}$ $= \frac{5}{4} \text{ o.e.}$	B1 M1 A1 (3)
(d)	$-1 < r < 1 \quad (\text{or } r < 1)$	B1 (1) (11)

8.

(a) $50\,000r^{n-1}$ (or equiv.) (Allow ar^{n-1} if $50\,000r^{n-1}$ is seen in (b))	B1	(1)
(b) $50\,000r^{n-1} > 200\,000$ (Using answer to (a), which must include r and n , and 200 000) (Allow equals sign or the wrong inequality sign) (Condone 'slips' such as omitting a zero)	M1	
$r^{n-1} > 4 \Rightarrow (n-1)\log r > \log 4$ (Introducing logs and dealing correctly with the power) (Allow equals sign or the wrong inequality sign)	M1	
$n > \frac{\log 4}{\log r} + 1$ (*)	Also	(3)
(c) $r = 1.09$: $n > \frac{\log 4}{\log 1.09} + 1$ or $n - 1 > \frac{\log 4}{\log 1.09}$ ($n > 17.086\dots$) (Allow equality)	M1	
Year 18 or 2023 (If one of these is correct, ignore the other)	A1	(2)
(d) $S_n = \frac{a(1-r^n)}{1-r} = \frac{50\,000(1-1.09^{10})}{1-1.09}$	M1 A1	
£760 000 (Must be this answer... nearest £10000)	A1	(3)
		9